

The early detection of prostate cancer in general practice:

Supporting patient choice

Brian R McAvoy MBChB MD FRNZCGP FRACGP FRCGP FRCP FACHAM BSc, Suzanne K Steginga PhD and Carole Pinnock PhD

On behalf of the Queensland Cancer Fund, the Australian Prostate Cancer Collaboration, the National Cancer Control Initiative, the Northern Section of the Urological Society of Australasia, the Prostate Cancer Foundation of Australia, The Cancer Council Victoria and the Royal Australian College of General Practitioners.

Correspondence to: Brian.McAvoy@cancervic.org.au

Background

Prostate cancer is the most commonly diagnosed cancer in men in New Zealand and the third most common cause of male cancer deaths, accounting for 3.8% of deaths.¹ The incidence of prostate cancer has risen rapidly over the past 15–20 years but this is likely to be the result of increased numbers of asymptomatic men undergoing prostate specific antigen (PSA) testing.²

Prostate cancer is very rare in men below the age of 50. About 90% of all new cases are in men aged 60 or older, and two-thirds of the men who die from the disease are over age 75.¹ The registration rate for new prostate cancers in Maori is lower than for non-Maori (71.3 per 100 000 population compared with 96.2 per 100 000 population), but these rates are based on relatively small numbers, and there has been variable recording of ethnic data, so the estimates for Maori men are less robust.²

Risk of diagnosis of, and deaths from, prostate cancer are very much age-related – the risk of diagnosis in men aged 70–79 being nearly seven times greater than that in men aged 50–59, and the risk of death being 21 times greater (Table 1). However, premature death from prostate cancer is more likely for

men diagnosed in their 50s compared to men in their 70s (Table 2). Family history increases the risk of developing prostate cancer – a man with a father or brother diagnosed has at least twice the risk of diagnosis,⁴ and this increases with decreasing age of the affected relative. There is also wide international variability in incidence rates of prostate cancer, with high rates in those of African American background and low rates in Japanese.⁵

Current recommendations

Population-based screening of asymptomatic men for prostate cancer in New Zealand is not recommended by the National Health Committee because of its lack of proven benefit and the potential harm arising from unnecessary radiotherapy, surgery or other treatment.⁶ This position is

also supported by the Cancer Council Australia, the Urological Society of Australasia, the Australian Prostate Cancer Collaboration, the Royal Australian College of General

Practitioners and the Australian Government Department of Health and Ageing. On the matter of opportunistic testing, these organisations and most clinical practice guidelines recommend that patients be fully informed of the risks, and limitations of the test benefits before making their own choice.^{7–11} The positive predictive value of combined abnormal PSA and digital rectal examination (DRE) is about 38–50%.^{13–14} For every hundred men who actually have prostate cancer, between 10% and 30% will have a normal PSA test result (up to 4.0ng/ml).^{15–16} DRE detects cancer in some men with PSA levels

below 4.0ng/ml.⁹ A graphic representation of what would happen to 1000 men who had a PSA test is shown in Figure 1. A man with a PSA level between 4.0 and 10.0ng/ml has an approximately 1-in-4 chance of having prostate

cancer. Non-cancer contributors to increased PSA include:¹⁷

- benign prostatic hypertrophy
- ejaculation in preceding 48 hours
- urinary tract infection

To make a fully informed choice about PSA testing men should also be aware that active treatments for prostate cancer such as surgery and radiotherapy... have significant potential for adverse outcomes

- urinary retention
- prostatitis
- prostatic massage but probably not routine DRE
- prostate needle biopsy.

To make a fully informed choice about PSA testing men should also be aware that active treatments for prostate cancer such as surgery and radiotherapy (external beam and brachytherapy) have significant potential for adverse outcomes which may include impotence, less commonly urinary incontinence, bowel problems and, although rarely, even death. The range of possible adverse outcomes following radical prostatectomy is shown in the text box in Figure 1.

However, these recommendations are controversial.¹⁸ The American Cancer Society and the American Urological Association recommend PSA and DRE screening beginning at age 50 in normal-risk men; in contrast, the American Medical Association and the American College of Physicians–American Society of Internal Medicine do not specifically recommend screening.¹⁹ Recently, the United States Preventive Services Task Force concluded that the net benefit of prostate cancer screening with PSA and DRE could not be determined and advise that screening for prostate cancer could not yet be either supported nor discounted.⁹ There is some emerging weak evidence for a benefit from screening^{20–21} but definitive evidence will not be available until the completion of current randomised controlled trials in the USA and Europe.

Supporting patient choice

Currently 25–43% of Australian men express interest in PSA testing and many undergo it.^{23–26} This may well reflect a greater recognition of patients' autonomy and a move away from a 'paternalistic' model of patient care.²⁷ It may also reflect increasing public awareness of prostate cancer and PSA testing – the Prostate Cancer Foundation of Australia (a patient support/advocacy

Table 1. Risk of diagnosis of, and death from, prostate cancer by age in Australia (per 1,000 men).

Age group	Of 1000 men at this age, number with a diagnosis of prostate cancer	Of 1000 men at this age, number who die from prostate cancer
0–49	1	<0.1
50–59	12	1
60–69	45	5
70–79	79	21
80–89	105	61
50–79	136	27

Data source: ABS and AIHW, based on 1997–2001 data³

Table 2. Premature death from prostate cancer

Of 1000 men diagnosed with prostate cancer, how many will die prematurely (before 80 years of age) from the disease?	
Age group (at diagnosis)	Number
50–59	600
60–69	500
70–79	380

Data source: Queensland Health, based on 1982–2000 data, with minimum two years follow-up³

group) has recently run a media campaign 'Be a man, see your doctor about prostate cancer'. The general practitioner (GP) has the task of responding to consumer demands for information in the face of conflicting viewpoints and uncertain medico-legal requirements.^{27,28}

Most patients' assessment of risk is primarily determined not by facts, but by emotions.²⁹ Decisions about health are strongly influenced by non-systematic thinking and involve such factors as framing effects, categorical thinking, low aversion, lay beliefs and memories of past experiences.^{30,31}

Because the evidence base is not clear-cut, a shared decision-making approach offers a way forward. This involves a two-way process and includes both medical and personal information. A framework for the competencies that medical practitioners need for shared decision making has been described:³²

1. Develop a partnership with the patient;

2. Establish or review the patient's preferences for information (such as amount or format);
3. Establish or review the patient's preferences for their role in decision-making (such as degree of involvement of self) and the existence and nature of any uncertainty about the course of action to take;
4. Ascertain and respond to the patient's ideas, concerns and expectations;
5. Identify choices (including ideas and information that the patient may have) and evaluate the research evidence in relation to the individual patient;
6. Present (or direct the patient to) evidence, taking into account competencies (2) and (3) and framing effects (how presentation of information may influence decision-making). Help the patient to reflect on, and assess, the impact of alternative decisions with regard to his or her lifestyle and values;

7. Make or negotiate a decision in partnership with the patient and resolve conflict;
8. Agree on an action plan and complete arrangements for follow-up.

GP education programme

Since GPs are the most likely source of information for PSA testing and subsequent referral, it is essential that they fully understand screening and treatment issues so that men in their care can make informed choices. With this in mind the Australian Prostate Cancer Collaboration partnered with the Queensland Cancer Fund, the Northern Section of the Urological Society of Australasia, the National Cancer Control Initiative (NCCI) and other groups to develop an educational programme and practice resource for GPs.

Based on materials produced by the Centers for Disease Control and Prevention in the United States,³³ resources were developed with input from GPs, public health practitioners, and academics, specialist urologists, consumer groups, psychologists, epidemiologists, a medicolegal expert and educationalists. The materials and workshops were piloted in Queensland and Victoria.

The education programme was designed to cover two main areas:

1. The medical context of screening, which includes risk and natural history of prostate cancer, potential benefits, and risks of screening for and treating prostate cancer, use and interpretation of PSA tests, and referral pathways for further investigation.
2. Shared decision-making, which covers medicolegal issues and informed consent responsibilities for opportunistic preventative health screening, understanding how men make health decisions, lay health beliefs, risk communication, and using a patient-centred approach for informed choice that elicits patients' values and priorities.

The education programme was designed as a 2.5 hour interactive

Figure 1. Outcome of PSA testing in 1000 men (From NZ Guidelines Group Booklet)




Figure 2. GP/patient show card

GP / Patient Show Card


1

The Early Detection of Prostate Cancer in General Practice: Supporting Patient Choice

These resource cards aim to support the general practitioner assisting a patient making a choice about prostate cancer testing. This decision ideally is both informed and consistent with the patient's personal preferences (1-4). Page 1 is a show card to use with your patients, giving suggestions for a discussion of pros and cons. Page 2 provides more detail for your use. Page 3 overviews PSA value ranges.

SIX DECISION STEPS – TALK TO YOUR PATIENT ABOUT:

1. What is your main concern?
2. What is prostate cancer and what tests are there?
3. What is your risk?
4. What are the pros and cons of early detection?
5. What is most important to you?
6. Your decision.



What is your risk? (5-6)

- Of 1000 men who are aged 50 years, 136 will be diagnosed with prostate cancer before the age of 80.
- Younger men have a smaller chance of a diagnosis than older men. But if they are diagnosed with prostate cancer, younger men are more likely to die prematurely from it. This is because there is more time for the cancer to progress and younger men are less likely to die of other causes.

What is the chance of a diagnosis of prostate cancer:

For a man in his 40s	1 in 1000
For a man in his 50s	12 in 1000
For a man in his 60s	45 in 1000
For a man in his 70s	79 in 1000

- Family history increases risk eg. a man with a father or brother diagnosed has at least twice the risk of a diagnosis. The risk increase is highest in relatives of men diagnosed before age 60 years, and decreases with increasing age of the affected relative.

The authors do not specifically recommend any treatment in this publication. Information on prostate disease is constantly being updated. We have made reasonable effort to ensure that information was current at the time of production (07/04/2005).


WHAT IS MOST IMPORTANT TO YOU?

For: Is this like you?

- ✓ I'm concerned that I might get prostate cancer.
- ✓ I want the best chance of finding it early, if I do get it.
- ✓ I'm not interested in waiting for all the proof to be in.
- ✓ I want to do everything possible to reduce my risk of dying from prostate cancer.

Against: Is this like you?

- ✓ I think my chance of getting prostate cancer is low.
- ✓ I am not convinced about the effectiveness of testing.
- ✓ I am more concerned about avoiding treatment side effects, if there's no guarantee I'd be reducing my risk of dying from prostate cancer.



SIX DECISION STEPS – TALK TO YOUR PATIENT ABOUT:

These steps apply to a patient concerned about or requesting a test for prostate cancer.

1. Clarify the patient's main concern

- General concern and/or:
- Lower urinary tract symptoms (LUTS) – reassurance that this does not make him more at risk. Refer to current NHMRC guidelines (2000) for LUTS management⁽⁷⁾
- Family history, confirm 1 or more 1st degree relatives diagnosed before 60 years.

2. Provide basic information on prostate cancer and tests available

- What the prostate is, where it is, that it grows bigger with age and can cause urinary symptoms over the age of 50 years.
- What prostate cancer is, how it is controlled by the male hormone, how in most men it grows quite slowly although rapidly in some.
- For early detection a blood test (PSA) and digital rectal exam (DRE) are needed. These are screening not diagnostic tests. If either or both of these are suspicious that does not necessarily indicate cancer. To find this out a prostate biopsy would then be needed.
- Chance of cancer given positive PSA test is 1 in 3. Chance of cancer given both abnormal PSA and DRE tests is 1 in 2. Cancer can still be present with a normal PSA.

3. Provide an estimate of this patient's risk of a diagnosis based on age and family history (assumes no previous PSA result available)

- Risk of getting prostate cancer increases with age. However given a diagnosis, older men are less likely to die prematurely from it - there is less time for the cancer to progress and more competing causes of death. Testing is not normally recommended in men with life expectancy <10 years.
- Of 1,000 50 year old men, about 136 will be diagnosed with prostate cancer, and 27 will die from prostate cancer, before the age of 80 years⁽⁸⁾. These risks increase with age. Less than one man in 1,000 will be diagnosed with prostate cancer in their 40s.
- These risks are population estimates, and assume that everyone is the same. Although we don't know what causes prostate cancer, international comparisons tell us that some men are at greater risk than others, possibly related to lifestyle and diet.
- Family history increases risk eg. a man with a father or brother diagnosed has at least twice the risk of a diagnosis⁽⁹⁾.
- Men who are between the ages of 50-75 years and men older than 45 years at increased risk (eg. family history)⁽⁹⁾ are most likely to benefit from the early detection of prostate cancer.

The authors do not specifically recommend any treatment in this publication. Information on prostate disease is constantly being updated. We have made reasonable effort to ensure that information was current at the time of production (07/04/2005).

4. Explain pros and cons of early detection

Pros

- Early prostate cancer has no symptoms - PSA testing can lead to the detection of prostate cancer before it causes symptoms and/or when it is still confined within the prostate gland (localised).
- Treatment for localised prostate cancer can potentially cure the disease.
- Prostate cancer that is still confined to the gland may progress over time.
- Prostate cancer that has spread beyond the prostate gland is usually no longer curable and treatment for advanced cancer has significant quality of life effects.

Cons

- Some prostate cancers grow slowly and don't threaten life, but detection and treatment for prostate cancer can affect quality of life.
- A PSA test can be abnormal when cancer is not present (happens 2/3 times for a positive test), however a biopsy is needed to find out. Explain what a biopsy involves.
- There is no clinical trial evidence yet that PSA testing programs save lives and whether men who are monitored by testing (screened) live longer. Because of this lack of evidence of effectiveness, medical authorities do not currently recommend population screening for prostate cancer, although this may change in the future.

Treatment Side Effects

- Potentially curative treatments for localised prostate cancer include surgery and radiation therapy (external beam and brachytherapy). These treatments are associated with significant risk of impotence, and less commonly urinary incontinence and bowel problems. Prevalence and profile of side effects vary for different treatment types⁽⁸⁻¹¹⁾.
- Advanced prostate cancer is treated primarily by hormonal manipulation and is associated with side effects such as impotence, loss of libido, fatigue, osteoporosis, and cognitive changes⁽¹²⁻¹³⁾.

5. Help the patient clarify their values

- Give examples of reasons men have given who have had or not had the test.
- Use table of "What is most important to you?" Ask the man to consider if any of these points seem like his feelings or views.

6. Confirm Decision

- Ask what questions he has. Check understanding.
- Does he want to decide now or take the written patient information and think about it?
- If the man chooses to be tested, discuss a prostate cancer risk management plan (see page 3).

For information about prostate cancer contact the Cancer Helpline on 13 11 20.

Useful Websites

• www.prostatehealth.org.au • www.urosoc.org.au • www.andrologyaustralia.org
• www.prostate.org.au

Figure 3. GP reference card

GP Reference Card

3



The Early Detection of Prostate Cancer in General Practice: Referral Guide for Prostate Testing

This information is provided with the view that if a man chooses to be tested, he hopes to have the cancer detected at an early stage, so that treatment options have the chance for cure. Testing of men with life expectancy less than 10 years is not normally recommended⁽¹⁴⁻¹⁵⁾. If PSA testing is performed a DRE is also recommended.

Normal ranges for PSA

Standard PSA normal range cut off: 4.0 ng/ml
Age-based normal ranges for PSA (ng/ml) Oesterling 1995⁽¹⁶⁾

Age range	50th percentile (median)	95th Percentile (upper limit of normal)
40-49	0.65	2.0
50-59	0.85	3.0
60-69	1.39	4.0
70-79	1.64	5.5

Differences in type of PSA assay can cause differences in age-based ranges

Men whose PSA is above the 50th (median) but below the 95th percentile have been shown to be at higher long term risk of prostate cancer compared with those below the median^(17,18).

Normal rate of change PSA has been reported to increase by about 3.3% per year (0.04 ng/ml at 60 yrs)⁽¹⁶⁾. A rate of change is observed from at least 3 PSA measurements over 12-18 months with a higher rate suggestive of increased cancer risk⁽¹⁹⁾.

Percentage free PSA (Free to Total Percentage or FTP) is lower when cancer is present and may be helpful to distinguish cancer from benign prostatic enlargement in men with intermediate total PSA ranges (2.0-10.0 ng/ml)⁽¹⁶⁾. Cancer is likely if FTP is below 10% and a low risk if FTP is over 25%.

Accuracy of test

The positive predictive value (chance of cancer given abnormal result) is about 30%^(15,20). The positive predictive value of combined abnormal PSA and DRE is about 38-50%^(21,22). For every hundred men who actually have prostate cancer, between 10% and 30% will have a normal PSA test result (up to 4.0 ng/ml)^(23,24). DRE detects cancer in some men with PSA levels below 4.0 ng/ml⁽¹⁵⁾.

Non cancer contributors to increases in PSA⁽²⁵⁾

1. Benign prostate enlargement – accounted for to some extent by using age-based reference ranges and percentage free-PSA (see left).
2. Ejaculation: both total PSA and % free PSA increase (can remain altered for 6-48 hours).
3. Urinary infection.
4. Urinary retention (48 hours after resolution, PSA decreased by 50%).
5. Prostatitis or sub-clinical prostate inflammation (can remain higher for at least 6 weeks following resolution).
6. Prostatic massage but probably not routine DRE (prudent to take blood prior to DRE).
7. Prostate needle biopsy.
8. Bicycle riding has been reported not to change the PSA level^(26,27).
9. Different manufacturer assays may cause variation (up to 10%).

* Other investigations to consider: MSU, Electrolytes, Creatinine

Consider referral if:

- PSA exceeds 4ng/ml or upper limit of normal for age range (95th percentile-see table)
- PSA rate of change from a normal base is high
- DRE indicates nodularity or hard prostate

Consider follow-up if:

- PSA is in upper ranges of normal for age (exceeds median)
- Patient has a family history of prostate cancer
- Patient requests testing for the purpose of early detection

Recommended follow-up intervals for the detection of early stage cancer may vary depending on the result of the PSA test^(20,28,29). Medicare Benefits Schedule for PSA as of November 2004 one patient episode in a 12 month period: refer to www.health.gov.au/pubs/mbbs.

The authors do not specifically recommend any treatment in this publication. Information on prostate disease is constantly being updated. We have made reasonable effort to ensure that information was current at the time of production (07/04/2005).

References



The Early Detection of Prostate Cancer in General Practice: Referral Guide for Prostate Testing

References

- Bird, S. Discussing benefits and risk with patients PSA testing. Australian Family Physician 2004. 33:266-267.
- Chan, E.C.Y. Sulmay, D.P. What men should know about Prostate-specific Antigen Screening before giving informed consent? American Journal of Medicine 1998. 105:266-274.
- Dunn, I.B. Kirk, D. Legal pitfalls in the diagnosis of prostate cancer. British Journal of Urology International 2000. 86:304-307.
- National Health and Medical Research Council 2004. General Guidelines for Medical Practitioners in Providing Information to Patients, Canberra. www.health.gov.au/nhmrc/publications/synopses/e57syn.htm
- Baade, P. Steginga, S.K. Pinnock, C. & Aitken, J. Communicating prostate cancer risk: what should we be telling our patients? Medical Journal of Australia. 2005. In Press.
- Johns, L.E. and Houston, R.S. A systematic review and meta-analysis of familial prostate cancer risk. British Journal of Urology International 2003. 91(9): 789-794.
- National Health and Medical Research Council. Clinical Practice Guidelines: The management of uncomplicated lower urinary tract symptoms in men. 2000 <http://www.health.gov.au/nhmrc/publications/synopses/cp42syn.htm>
- US Preventive Services Task Force. Screening for prostate cancer: recommendation and rationale. Annals of Internal Medicine 2002. 137(11): 915-6.
- National Health and Medical Research Council and Australian Cancer Network. Clinical Practice Guidelines: Evidence-based information and recommendations for the management of localised prostate cancer 2003. Commonwealth of Australia, Canberra.
- Steginga, S.K. Occhipinti, S. Gardiner, R. A. Heathcote, P. & Yaxley, J.A. Prospective Study of Men's Psychological and Decision-Related Adjustment after Treatment for Localised Prostate Cancer. Urology 2004. 63: 751-756.
- Steinbeck, G. Helgesen, F. Adolfsson, J. et al. Quality of life after radical prostatectomy or watchful waiting. New England Journal of Medicine 2002. 347(11): 790-6.
- Holzbeler, J.E. Castle and J. Thrasher Complications of androgen deprivation therapy: prevention and treatment. Oncology Huntington 2004. 18(3): 303-9.
- Green, H. Pakenham, K.I. Headley, B. et al. Quality of life compared during pharmacological treatments and clinical monitoring for non-localised prostate cancer: a randomised controlled trial. British Journal of Urology International 2004. 93(7): 975-9.
- Oberg, A. Allan, J. Frame, P. Homer C. et al. Screening for prostate cancer: recommendation and rationale. US Preventive Services Task Force. Annals of Internal Medicine 2002. 137(11):915-6.
- Harris, R. and Lohr, K.N. Screening for prostate cancer: an update of the evidence for the U.S. Preventive Services Task Force. Annals of Internal Medicine 2002. 137(11): 917-29.
- Oesterling, J.E. Jacobsen, S.J. Klee, G.G. et al. Free, complexed and total serum prostate specific antigen: the establishment of appropriate reference ranges for their concentrations and ratios. Journal of Urology 1995. 154(3):1090-5.
- Fang, J. Metter, E.J. Landis, P. et al. Low levels of prostate-specific antigen predict long-term risk of prostate cancer: results from the Baltimore Longitudinal Study of Aging. Urology 2001. 58(3):411-6.
- Gann, P.H. Hennekens, C.H. Stampfer, M.J. A prospective evaluation of plasma prostate-specific antigen for detection of prostatic cancer. Journal of the American Medical Association 1995. 273(4):289-94.
- Carter, H.B. Pearson, J.D. Metter, E.J. et al. Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease. cancer. Journal of the American Medical Association 1992. 267(16):2215-20.
- Hugosson, J. Aus, G. Lilja, H. et al. Prostate specific antigen based biennial screening is sufficient to detect almost all prostate cancers while still curable. Journal of Urology 2003. 169(5):1694-6.
- Mistry, K. and Cable, G. Meta-analysis of prostate-specific antigen and digital rectal examination as screening tests for prostate carcinoma. Journal of the American Board of Family Practice 2003. 16(2): 95-101.
- Coley, C.M. et al. Early detection of prostate cancer. Part 1: Prior probability and effectiveness of tests. The American College of Physicians. Annals of Internal Medicine. 1997. 126(5):394-406.
- Narain, V.M.L. Cher, et al. 2002. Prostate cancer diagnosis, staging and survival. Cancer and Metastasis Reviews 21: 17-27.
- Schroder, F.J. van der Cruisen-Koeter, et al. 2000. Prostate cancer detection at low prostate specific antigen. Journal of Urology 163(3): 806-812.
- Polascik, T.J. Oesterling, J.E. Partin, A.W. Prostate specific antigen: a decade of discovery - what we have learned and where we are going. Journal of Urology. 1999. 162(2):293-306.
- Crawford, E.D. 3rd, Mackenzie, S.H. Safford, H.R. Capriola, M. The effect of bicycle riding on serum prostate specific antigen levels. Journal of Urology. 1996. 156(1):103-5.
- Lubold, H.J. Peck, K.D. Oberpenning, F. et al. Bicycle riding has no important impact on total and free prostate-specific antigen serum levels in older men. Urology. 2003. 61(6):1177-80.
- Draisma, G. Boer, R. Otto, S.J. et al. Lead times and overdiagnosis due to prostate-specific antigen screening: estimates from the European Randomized Study of Screening for Prostate Cancer. Journal of the National Cancer Institute 2003. 95(12):838-9.
- Carter, H.B. Epstein, J.I. Chan, D.W. et al. Recommended prostate-specific antigen testing intervals for the detection of curable prostate cancer. Journal of the American Medical Association 1997. 277(18):1456-60.

The authors do not specifically recommend any treatment in this publication. Information on prostate disease is constantly being updated. We have made reasonable effort to ensure that information was current at the time of production (07/04/2005).

workshop and has been accredited for continuing professional development points under the Royal Australian College of General Practitioners' Quality Assurance and Continuing Professional Development Program and the Australian College of Rural and Remote Medicine's Professional Development Program. The workshop involves presentations by a urologist, a general practitioner and a communication/decision support expert. Workshop materials developed include a set of PowerPoint slides and notes for the presenters and three case studies for small group discussion.

In addition, practice resources were developed that included an interactive decision aid and brief evidence-based summary card for use in the consultation. The interactive decision aid is a GP/patient show card (Figure 2) with an anatomical diagram of the male pelvis, six decision steps, current age based risk estimates for prostate cancer and a values clarification exercise. The summary card also contains more detail for the informed choice discussion and a GP reference card (Figure 3) with a referral guide covering age-based PSA reference ranges and non-cancer causes of elevated PSA readings. The resources were designed so as to be easily scanned visually and suitable for use in a brief consultation. With financial support from The Cancer Council Australia and the National

Seniors Foundation, these materials have been distributed to all GPs in Australia with a recent edition of the *Australian Family Physician*. They are also downloadable as a pdf on the NCCI's website – www.ncci.org.au.

Evaluation of the education programme using a single arm pre-post test design has shown high satisfaction with the workshop and resource materials. Participants have shown significantly increased knowledge, level of understanding and confidence in discussing testing with asymptomatic men and initiating discussions about testing with patients.³²

Since completion of the pilot workshops, over 450 GPs in Queensland alone and 121 in Victoria have attended the programme with further workshops scheduled in other states. A 'train-the-trainer' programme has been developed by the NCCI, and is due to commence in New South Wales in 2006.

Conclusion

In a health topic characterised by divergent viewpoints, this educational programme provides evidence of the benefits of taking a collaborative and consultative approach, and closely linking programme development to GPs' expressed needs. Focusing the educational programme on shared decision-making appears to be effective in improving GPs' knowledge, level of understanding and confidence.

Acknowledgement

We gratefully acknowledge the assistance of Dr Peter Heathcote, Dr John Preston Mr Ross Snow, Mr Michael Chamberlain and Associate Professor Damien Bolton in the development and piloting of this education programme. We also gratefully acknowledge the support of the Queensland Cancer Fund, the Australian Prostate Cancer Collaboration, the Prostate Cancer Foundation of Australia, the Northern section of the Urological Society of Australasia, The Cancer Council Victoria, the National Cancer Control Initiative, the Brisbane North Division of General Practice, the Northern Division of General Practice, Inner Eastern Melbourne Division of General Practice, Melbourne Division of General Practice, Greater South Eastern Division of General Practice and Dandenong District Division of General Practice.

New Zealand GPs interested in piloting a local version of the workshop should contact:

Dr Quinten King, Consultant Urologist
Palmerston North Hospital
Ruahine Street, Palmerston North

Tel: (06) 356 9169

Email: Quinten.King@midcentral.co.nz

Competing interests

None declared

References

1. Durham J. Population screening for prostate cancer. A systematic review. Wellington, New Zealand: New Zealand Guidelines Group; 2002.
2. New Zealand Health Information Service. Cancer: New Registrations and Deaths 1998. Wellington, New Zealand: New Zealand Health Information Service; 2002:124.
3. Baade PD, Steginga SK, Pinnock CB, et al. Communicating prostate cancer risk: what should we be telling our patients? *Med J Aust* 2005; 182:472-475.
4. Johns LE, Houlston RS. A systematic review and meta-analysis of familial prostate cancer risk. *B J U Int* 2003; 91:789-794.
5. Parkin DM, Whelan SL, Ferlay J, et al, editors. Cancer incidence in five continents. Volume VIII. IARC Scientific Publications No. 155. Lyon, France: International Agency for Research on Cancer (WHO) and International Association of Cancer Registries, 2002.
6. National Health Committee. Prostate Cancer Screening in New Zealand. Report to the Minister of Health. Wellington, New Zealand: National Health Committee; 2004.
7. Australian Health Technology Advisory Committee. Prostate Cancer Screening. Canberra: Australian Government Publishing Service; 1996.
8. Coley CM, Barry MJ, Mulley AG. Clinical guidelines part III: Screening for prostate cancer. *Ann Intern Med* 1997; 126: 480-484.
9. Harris R, Lohr KN. Screening for prostate cancer: an update of the evidence for the US Preventive Services Task Force. *Ann Intern Med* 2005; 137:917-929.
10. Talcott JA. What patients should be told before agreeing to a blood test that could change their lives. *Urology* 2003; 61:7-9.
11. Wilkinson S, Chodak G. Informed consent for prostate specific antigen screening. *Urology* 2003; 61:2-4.

12. Hugosson J, Aus G, Lilja H, et al. Prostate specific antigen based biennial screening is sufficient to detect almost all prostate cancers while still curable. *J Urology* 2003; 169:1694-1696.
13. Mistry K, Cable G. Meta-analysis of prostate specific antigen and digital rectal examination as screening tests for prostate carcinoma. *J Am Board Fam Pract* 2003; 16:95-101.
14. Coley CM, Barry MJ, Fleming C, et al. Early detection of prostate cancer. Part 1: Prior probability and effectiveness of tests. *The American College of Physicians. Ann Int Med* 1997; 126:394-406.
15. Narain VML, Cher ML, Wood DP Jr. Prostate cancer diagnosis, staging and survival. *Cancer and Metastasis Reviews* 2002; 21: 17-27.
16. Schroder FH, van der Crujisen-Koeter I, de Koning HJ, et al. Prostate cancer detection at low prostate specific antigen. *J Urol* 2000; 163:806-812.
17. Polascik TJ, Oesterling JE, Partin AW. Prostate specific antigen: a decade of discovery - what we have learned and where we are going. *J Urol* 1999; 162:293-306.
18. Catalona WJ. Informed consent for prostate specific antigen screening. *Urology* 2003; 61:17-19.
19. Zoorob R, Anderson R, Cefalu C, et al. Cancer screening guidelines. *Am Fam Physician* 2001; 63:1101-1112.
20. Baade PD, Coory MD, Aitken JF. International trends in prostate cancer mortality: the decrease is continuing and spreading. *Cancer Causes Control* 2004; 15:237-241.
21. Bill-Axelsson A, Holmberg L, Ruoto M, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *New Engl J Med* 2005; 352: 1977-1984.
22. Johansson J, Andren O, Andersson S, et al. National risk and history of early, localised prostate cancer. *JAMA* 2004; 291:2713-2719.
23. Slevin TJ, Donnelly N, Clarkson JP, English DR, Ward JE. Prostate cancer testing: behaviour, motivation and attitudes among Western Australian men. *Med J Aust* 1999; 171:185-8.
24. Smith DP, Armstrong BK. Prostate specific antigen testing in Australian and association with prostate cancer incidence in New South Wales. *Med J Aust* 1998; 169:17-20.
25. Steginga SK, Occhipinti S, McCaffrey J, Dunn J. Men's attitudes to prostate cancer and seeking prostate specific antigen testing. *J Cancer Educ* 2001; 16:42-5.
26. Weller D, Pinnock C, Silagy C, Hiller JE, Marshall VR. Prostate cancer testing in SA men: influence of sociodemographic factors, health beliefs and LUTS. *Aust NZ J Public Health* 1998; 22:400-2.
27. Bird S. Risk management. Discussing benefits and risks with patients. PSA testing. *Aust Fam Physician* 2004; 33:266-267.
28. Dunn IB, Kirk D. Legal pitfalls in the diagnosis of prostate cancer. *B J U Int* 2000; 86:304-307.
29. Paling J. Strategies to help patients understand risk. *BMJ* 2003; 327:745-748.
30. Steginga SK, Occhipinti S, Gardiner RA, et al. Making Decisions about Treatment for Localised Prostate Cancer. *Brit J Urol Int* 2002; 89:255-260.
31. Redelmeier DA, Rozin P, Kahneman D. Understanding patients' decisions: Cognitive and emotional perspectives. *JAMA* 1993; 270:72-76.
32. Toule A, Gadolphin W. Framework for teaching and learning informed shared decision making. *BMJ* 1999; 319:766-771.
33. Harris RP, Flood AB, Berge V, et al. Screening for prostate cancer: Sharing the Decision Slide set. On line: National Center for Chronic Disease Prevention and Health Promotion; (updated 2005 January 20; cited 2005 May 20), available from: http://www.cdc.gov/cancer/prostate/screening/slide_index.htm.
34. Steginga SK, Pinnock C, Baade PD et al. An educational workshop on the early detection of prostate cancer. *Aust Fam Physician* 2005; 34:889-890.